

Total decay width of $H \rightarrow gg$ using the infinite-order scale-setting approach based on intrinsic conformality*

Chu-Tian Gao(高楚天)^{1†} Xing-Gang Wu(吴兴刚)^{1‡} Xu-Dong Huang(黄旭东)^{2,3§} Jun Zeng(曾军)^{4¶}

¹Department of Physics, Chongqing Key Laboratory for Strongly Coupled Physics, Chongqing University, Chongqing 401331, China

²Institute of High Energy Physics, CAS, Beijing 100049, China

³School of Physics, University of Chinese Academy of Sciences, Beijing 100049, China

⁴INPAC, Key Laboratory for Particle Astrophysics and Cosmology (MOE), Shanghai Key Laboratory for Particle Physics and Cosmology, School of Physics and Astronomy, Shanghai Jiao Tong University, Shanghai 200240, China

Abstract: We conducted a detailed study on the properties of the total decay width of the Higgs decay channel $H \rightarrow gg$ up to α_s^6 -order QCD corrections by using the newly suggested infinite-order scale-setting approach, which is based on both the principle of maximum conformality and intrinsic conformality. This approach is called PMC_∞ . By using the PMC_∞ approach, we observed that the conventional renormalization scale ambiguity in perturbative QCD calculation is eliminated, and the residual scale dependence due to unknown higher-order terms can also be highly suppressed. We then obtained an accurate perturbative QCD prediction on the total decay width, e.g., $\Gamma(H \rightarrow gg)|_{\text{PMC}_\infty} = 336.42^{+7.01}_{-6.92}$ keV, where the errors are squared averages of those from all the mentioned error sources.

Keywords: pQCD, renormalization, Higgs decay

DOI: 10.1088/1674-1137/ac92da

In quantum chromodynamics (QCD), the Higgs boson plays an important role in the precision test of the Standard Model (SM), and it is also helpful for searching new physics beyond the SM. The Higgs boson decays into two gluons. This is an important channel for studying the Higgs phenomenology [1]. The coupling of the Higgs to gluons is predominantly mediated by the top quark within the SM, and the high-order QCD corrections to this process can be evaluated by means of an effective theory in which the top quark is integrated out [2]. At present, the perturbative QCD (pQCD) correction to the total decay width of the Higgs decay channel $H \rightarrow gg$, e.g., $\Gamma(H \rightarrow gg)$, is calculated up to next-to-next-to-next-to-next-to-leading order ($N^4\text{LO}$) in the limit of a large top-quark mass [3–11]. Therefore, we face the opportunity of achieving precise pQCD prediction on $\Gamma(H \rightarrow gg)$.

It is helpful to reduce the pQCD uncertainties as much as possible. Among them, the error caused by us-

ing the conventional scale-setting approach is usually treated as an important systematic error for pQCD prediction. Such an error in making fixed-order prediction occurs because one conventionally assumes an arbitrary renormalization scale to perform the numerical analysis, which is usually chosen as the typical momentum flow of the process, the one assumed to be the effective virtuality of the strong interaction, the one that eliminates large logs to achieve a more convergent series, etc. This *ad hoc* assignment of renormalization scale causes mismatching of α_s and the corresponding coefficients. Thus, the coefficients of the QCD running coupling at each order strongly depend on the choice of renormalization scale as well as the renormalization scheme. However, as indicated by the renormalization group invariance, a physical observable must be independent of the choice of renormalization scale. In the literature, the principle of maximum conformality (PMC) [12–15] has been suggested to

Received 2 July 2022; Accepted 19 September 2022; Published online 28 September 2022

* Supported by the Chongqing Graduate Research and Innovation Foundation (CYB21045, ydstd1912), the Natural Science Foundation of China (12175025, 12147102), and the Fundamental Research Funds for the Central Universities (2020CQJQY-Z003, 2021CDJZYJH-003)

† E-mail: gaoct@cqu.edu.cn

‡ E-mail: wuxg@cqu.edu.cn

§ E-mail: hxud@cqu.edu.cn

¶ E-mail: zengj@cqu.edu.cn



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Article funded by SCOAP³ and published under licence by Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd

remove such renormalization scale ambiguity. It is well known that the α_s -running behavior is governed by the renormalization group equation (RGE). Then, the $\{\beta_i\}$ -terms that emerged in the pQCD series can be inversely adopted for fixing the correct α_s -value of a high-energy process. The purpose of PMC is to rightly determine the effective coupling constant of the process (whose argument is called the PMC scale) with the help of the RGE [16, 17], whose prediction is found to be independent of any choice of renormalization scale and satisfies the requirement of renormalization group invariance. The PMC scale-setting procedure agrees with the standard scale-setting procedure of Gell-Mann and Low [18] in the QED Abelian limit (small number of colors, $N_C \rightarrow 0$ [19]).

Many successful PMC applications have been explored in the literature. Previously, the PMC has been applied for dealing with the decay width $\Gamma(H \rightarrow gg)$ [20–22]. Note that the PMC was originally introduced as a multi-scale approach, in which distinct effective couplings (and hence the PMC scales) at each order were derived because different categories of $\{\beta_i\}$ -terms occur at each order. Furthermore, because the same category of $\{\beta_i\}$ -terms emerges at different orders, the determined PMC scales are in perturbative form. This leads to the fact that the precision of the PMC scales at higher orders decreases with the increment of perturbative orders, given that fewer $\{\beta_i\}$ -terms are known for fixing the value of higher-order α_s . Thus, the PMC multi-scale approach shall have explicit residual scale dependence [23], and if the convergence of the perturbative series of the PMC scale is weak, such residual scale dependence could be large [24].

By further exploiting the intrinsic conformality (iCF) property into PMC, a new infinite-order scale-setting approach, called the PMC_∞ approach, has been recently proposed in Ref. [25]. The PMC_∞ approach follows from the PMC, and its resultant conformal coefficients are the same as those of PMC at each perturbative order but sets the effective PMC scales at each order by requiring all the scale-dependent $\{\beta_i\}$ -terms at each order to vanish exactly and separately [25]. Following this approach, the newly fixed PMC scales at each order are in definite form and are no longer in perturbative series. Thus, the residual scale dependence of the previous PMC scales, owing to their previous perturbative nature, can be exactly eliminated. This indicates that the precision of the previous PMC predictions on the total decay width $\Gamma(H \rightarrow gg)$

[20–22] may be further improved by applying the PMC_∞ approach. It is thus interesting to conduct a detailed study on $\Gamma(H \rightarrow gg)$ by using the PMC_∞ approach.

Practically, the width of the Higgs decays into two gluons at the α_s^6 -order level can be expressed as

$$\Gamma(H \rightarrow gg) = \frac{M_H^3 G_F}{36 \sqrt{2} \pi} \left[\sum_{k=0}^4 C_k(\mu_r) \alpha_s^{k+2}(\mu_r) \right], \quad (1)$$

where the Fermi constant $G_F = 1.16638 \times 10^{-5} \text{ GeV}^{-2}$, $a_s = \alpha_s/4\pi$, and μ_r stands for an arbitrary renormalization scale. The perturbative coefficients $C_{k \in [0,4]}(\mu_r)$ at the initial scale of $\mu_r = M_H$ under conventional $\overline{\text{MS}}$ -scheme can be read from Refs. [2–11]. As has been argued in Refs. [20–22], it is important to first transform them into those under a physical momentum space subtraction scheme (mMOM-scheme) [26–31] to avoid the ambiguities of fixing the PMC scales with the help of the RGE. The mMOM-scheme is gauge dependent, a detailed discussion of gauge dependence after applying the PMC has been done in Ref. [32], which shows that if the gauge parameter $\xi \in [-1, 1]$, the mMOM prediction will have weaker ξ -dependence. Concerning definiteness, we adopt the Landau gauge ($\xi = 0$) to conduct the analysis, whose corresponding coefficients $C_k(\mu_r)$ at any renormalization scale μ_r can be achieved by recursively applying the RGE. The explicit expressions for the required coefficients up to α_s^6 -order level can be found in Refs. [21, 22].

Owing to the iCF property, we can divide the N^4LO -level total decay width into five conformal subsets,

$$\Gamma(H \rightarrow gg) = \frac{M_H^3 G_F}{36 \sqrt{2} \pi} \sum_{n=1}^V \Gamma_n, \quad (2)$$

which collect together the same category of non-conformal terms into each subset and ensure the scheme independence of each subset via the commensurate scale relations among different orders [33]. Each conformal subset satisfies the scale invariant condition,

$$\left(\mu_r^2 \frac{\partial}{\partial \mu_r^2} + \beta(\alpha_s) \frac{\partial}{\partial \alpha_s} \right) \Gamma_n = 0. \quad (3)$$

More explicitly, we have

$$\Gamma_{\text{I}} = A_{\text{Conf}} \left[a_s^2(\mu_r) + 2B_{\beta_0} \beta_0 a_s^3(\mu_r) + (3B_{\beta_0}^2 \beta_0^2 + 2B_{\beta_0} \beta_1) a_s^4(\mu_r) + (7B_{\beta_0}^2 \beta_1 \beta_0 + 4B_{\beta_0}^3 \beta_0^3 + 2B_{\beta_0} \beta_2) a_s^5(\mu_r) + (8B_{\beta_0}^2 \beta_2 \beta_0 + \frac{47}{3} B_{\beta_0}^3 \beta_1 \beta_0^2 + 5B_{\beta_0}^4 \beta_0^4 + 2B_{\beta_0} \beta_3 + 4B_{\beta_0}^2 \beta_1^2) a_s^6(\mu_r) \right], \quad (4)$$

$$\Gamma_{\text{II}} = B_{\text{Conf}} \left[a_s^3(\mu_r) + 3C_{\beta_0} \beta_0 a_s^4(\mu_r) + (6C_{\beta_0}^2 \beta_0^2 + 3C_{\beta_0} \beta_1) a_s^5(\mu_r) + \left(\frac{27}{2} C_{\beta_0}^2 \beta_1 \beta_0 + 10C_{\beta_0}^3 \beta_0^3 + 3C_{\beta_0} \beta_2 \right) a_s^6(\mu_r) \right], \quad (5)$$

$$\Gamma_{\text{III}} = C_{\text{Conf}} \left[a_s^4(\mu_r) + 4D_{\beta_0} \beta_0 a_s^5(\mu_r) + (10D_{\beta_0}^2 \beta_0^2 + 4D_{\beta_0} \beta_1) a_s^6(\mu_r) \right], \quad (6)$$

$$\Gamma_{\text{IV}} = D_{\text{Conf}} \left[a_s^5(\mu_r) + 5E_{\beta_0} \beta_0 a_s^6(\mu_r) \right], \quad (7)$$

$$\Gamma_{\text{V}} = E_{\text{Conf}} \left[a_s^6(\mu_r) \right]. \quad (8)$$

Here, A_{Conf} , B_{Conf} , C_{Conf} , D_{Conf} , and E_{Conf} are conformal coefficients, and $B_{\beta_0} = \ln \mu_r^2 / \mu_1^2$, $C_{\beta_0} = \ln \mu_r^2 / \mu_{\text{II}}^2$, $D_{\beta_0} = \ln \mu_r^2 / \mu_{\text{III}}^2$, $E_{\beta_0} = \ln \mu_r^2 / \mu_{\text{IV}}^2$. The PMC_∞ scales are $\mu_{\text{I}, \dots, \text{IV}}$; they can be fixed by using the scale invariant condition (3). To match the mMOM-scheme perturbative series, the $\{\beta_i\}$ -functions under the mMOM-scheme should be adopted; their explicit forms up to five-loop level are available in Ref. [34]. Then, following the standard PMC_∞ scale-setting procedures, the conformal coefficients and PMC_∞ scales can be progressively derived from the known coefficients C_k via a step-by-step manner. For examples, we have $A_{\text{Conf}} = C_0$; the conformal coefficient B_{Conf} can be determined by setting $n_f = \frac{33}{2}$ ¹⁾ to drop off the β_0 terms in C_1 , and the PMC_∞ scale μ_{I} can be fixed by using the known conformal coefficients, i.e., A_{Conf} , B_{Conf} , the $\{\beta_0\}$ -terms of C_1 , etc. For convenience, we provide all the required conformal coefficients and PMC_∞ scales in the Appendix.

Then, we can transform the original perturbative series (1) into the following conformal series:

$$\Gamma(H \rightarrow gg) = \frac{M_H^3 G_F}{36 \sqrt{2} \pi} \left[A_{\text{Conf}} a_s^2(\mu_{\text{I}}) + B_{\text{Conf}} a_s^3(\mu_{\text{II}}) + C_{\text{Conf}} a_s^4(\mu_{\text{III}}) + D_{\text{Conf}} a_s^5(\mu_{\text{IV}}) + E_{\text{Conf}} a_s^6(\mu_{\text{V}}) \right]. \quad (9)$$

The PMC_∞ scales are definite and have no perturbative nature. Thus, they exactly avoid the residual scale ambiguity due to unknown higher-order terms in the perturbative series of the original PMC scales. Numerically, the first four PMC_∞ scales are

$$\{\mu_{\text{I}}, \mu_{\text{II}}, \mu_{\text{III}}, \mu_{\text{IV}}\} = \{50.1, 46.0, 63.0, 61.3\} (\text{GeV}), \quad (10)$$

which are invariant to any choice of renormalization scale and avoid the conventional renormalization scale ambiguity. Note that these PMC_∞ scales are around

$M_H \exp(-5/6) \sim 54$ GeV, which is suggested by the Gell-Mann Low scheme [18], in which $\exp(-5/6)$ is a result of the convention that defines the minimal dimensional regularization scheme. At present, the PMC_∞ scale μ_{V} at the highest order cannot be determined, since there is no $\{\beta_i\}$ -terms to fix its magnitude. As usual, we adopt $\mu_{\text{V}} = \mu_{\text{IV}}$ [15], which ensures the scheme independence of the resultant conformal series. Numerically, we found that due to the coefficient E_{Conf} is free of divergent renormalon terms, the magnitude of the final term is negligibly small, and the uncertainty of the total decay width caused by different choice of μ_{V} is negligible.

To do the numerical calculation, we set the top-quark pole mass $M_t = 172.5 \pm 0.7$ GeV, and $M_H = 125.25 \pm 0.17$ GeV [35]. The QCD asymptotic scale Λ can be determined by using the world average of α_s at the M_Z scale, e.g., $\alpha_s(M_Z) = 0.1179 \pm 0.0009$ [35]. As a subtle point, note that we need to transform the asymptotic scale from the $\overline{\text{MS}}$ -scheme to the mMOM-scheme by using the Celmaster-Gonsalves relation [26–29].

By setting all input parameters to their central values, we first obtained the decay width $\Gamma(H \rightarrow gg)$ up to different α_s -orders under conventional (Conv.) and PMC_∞ scale-setting approaches, as shown in Fig. 1. At the $O(\alpha_s^2)$ -order level, the perturbative series of $\Gamma(H \rightarrow gg)$ does not have $\{\beta_i\}$ -terms to fix μ_{I} , and the PMC_∞ and conventional predictions are the same and scale dependent. Figure 1 shows that the decay width $\Gamma(H \rightarrow gg)$ under conventional scale-setting approach has a strong dependence on μ_r , which becomes progressively smaller as more loop terms are included. Figure 1 also shows that the decay width $\Gamma(H \rightarrow gg)$ at $O(\alpha_s^3)$ -order and higher orders under PMC_∞ scale-setting are independent of any choice of renormalization scale owing to the fact that the scale-dependent nonconformal terms have been exactly eliminated.

Next, we present the decay width $\Gamma(H \rightarrow gg)$ up to different loop QCD corrections under conventional and PMC_∞ scale-setting approaches in Table 1. To show the perturbative property, we define a ratio

$$\kappa_n = \left| \frac{\Gamma^{O(\alpha_s^{n+2})} - \Gamma^{O(\alpha_s^{n+1})}}{\Gamma^{O(\alpha_s^{n+1})}} \right|, \quad (11)$$

which indicates how the "known" prediction $\Gamma^{O(\alpha_s^{n+1})}$ is affected by the one-order-higher terms. As for the PMC_∞ series, we have $\kappa_1 > \kappa_2 > \kappa_3 > \kappa_4$ for any choice of μ_r , indicating that the relative difference between the two nearby orders becomes smaller when more loop terms are included. This feature is consistent with the perturbative nature of the series and indicates that one can obtain more precise predictions by including more loop terms. Con-

1) Due to the $\beta_0 = 11 - \frac{2}{3}n_f$, to remove the β_0 -dependent terms from the coefficients C_i is equivalent to set $n_f = \frac{33}{2}$.




Fig. 1. (color online) Decay width $\Gamma(H \rightarrow gg)$ under conventional and PMC scale-setting approaches, respectively. The solid line with big dot, dash-dot line, dotted line, dashed line, and solid line are predictions up to $\mathcal{O}(\alpha_s^2)$, $\mathcal{O}(\alpha_s^3)$, $\mathcal{O}(\alpha_s^4)$, $\mathcal{O}(\alpha_s^5)$, and $\mathcal{O}(\alpha_s^6)$, respectively.

cerning the conventional series, as shown in Fig. 2, there are crossovers for $\kappa_{2,3,4}$ within the range of $\mu_r \in [M_H/2, 2M_H]$, and the ratios vary from 0 to 13%, 4.8%, and 1.0% for κ_2 , κ_3 , and κ_4 , respectively.

Moreover, to show the convergence of the perturbative series explicitly, we present the magnitude of each loop term for the four-loop approximants $\Gamma^{\mathcal{O}(\alpha_s^4)}$ in Table 2, which shows that the relative importance of the LO-terms: NLO-terms: N²LO-terms: N³LO-terms: N⁴LO-terms for the conventional series is

$$1 : +53.3_{-16.7}^{+13.5}\% : +6.6_{-16.4}^{+15.7}\% : -4.0_{-2.1}^{+6.9}\% : -1.6_{-2.7}^{+0.6}\%,$$

where the central values are those for $\mu_r = M_H$, and the errors are those for $\mu_r \in [\frac{1}{2}M_H, 2M_H]$. The scale depend-




Fig. 2. (color online) Results for the ratio κ_n versus renormalization scale μ_r under conventional scale-setting approach for different loop corrections. The ratios under PMC $_{\infty}$ approach are scale invariant.

ence for each loop term is large. However, owing to the cancellation of scale dependence among different orders, the net scale dependence is small, e.g., $(\begin{smallmatrix} +0.6\% \\ -0.3\% \end{smallmatrix})$ for $\mu_r \in [\frac{1}{2}M_H, 2M_H]$. In contrast, there is no renormalization scale dependence for each loop term of the PMC $_{\infty}$ prediction $\Gamma^{\mathcal{O}(\alpha_s^4)}$. More explicitly, we also present the value of each loop-term (LO, NLO, N²LO, N³LO, or N⁴LO) for $\Gamma^{\mathcal{O}(\alpha_s^4)}$ under the PMC $_{\infty}$ approach in Table 2. At the four-loop level, the PMC $_{\infty}$ series already represents good convergent behavior, and the relative importance of the LO-terms: NLO-terms: N²LO-terms: N³LO-terms: N⁴LO-terms becomes

$$1 : +31.8\% : -14.0\% : -6.5\% : +1.0\%,$$

whose magnitudes are scale invariant, indicating that the PMC $_{\infty}$ perturbative series represents the intrinsic perturbative behavior of $\Gamma(H \rightarrow gg)$. For comparison, we also show the numerical results under the PMC multi-scale approach (PMCm) in Table 2, which still has some residual scale-dependence. However, its numerical effect is smaller than that of the conventional approach. Detailed formulas for the PMCm approach can be found in Ref. [22].

Now, after eliminating the renormalization scale ambiguities, there are still some other error sources for the pQCD prediction of the total decay width, such as the α_s fixed-point error $\Delta\alpha_s(M_Z)$, the Higgs mass uncertainty ΔM_H , and the top-quark pole mass uncertainty ΔM_t . Up to α_s^6 -order, we have

Table 1. Results for the decay width $\Gamma(H \rightarrow gg)$ (unit: keV) and κ_n for different loop corrections under conventional and PMC $_{\infty}$ scale-setting approaches, respectively. The NLO and higher order PMC $_{\infty}$ predictions are scale independent, while, under the conventional scale-setting approach, the central values are those for $\mu_r = M_H$, and the errors are those for $\mu_r \in [M_H/2, 2M_H]$.

	$n=2$	$n=3$	$n=4$	$n=5$	$n=6$	κ_1	κ_2	κ_3	κ_4
$\Gamma^{\mathcal{O}(\alpha_s^4)} _{\text{Conv.}}$	$219.86_{-39.50}^{+54.05}$	$335.46_{-38.59}^{+41.34}$	$349.71_{-13.68}^{+2.52}$	$340.95_{-7.67}^{+1.00}$	$337.45_{-1.18}^{+1.94}$	[38%, 65%]	[0, 13%]	[0, 4.8%]	[0, 1.0%]
$\Gamma^{\mathcal{O}(\alpha_s^4)} _{\text{PMC}_{\infty}}$	$219.86_{-39.50}^{+54.05}$	389.86	342.09	334.05	336.42	[53%, 95%]	12%	2.4%	0.7%

Table 2. Values (unit: keV) of each loop-term (LO, NLO, N²LO, and N³LO) for the four-loop prediction $\Gamma^{O(\alpha_s^6)}$ under the conventional, PMC_∞ scale-setting, and PMCm scale-setting approaches, respectively. The PMC_∞ predictions are scale independent, while, under the conventional and PMCm scale-setting approaches, the central values are those for $\mu_r = M_H$, and the errors are those for $\mu_r \in [M_H/2, 2M_H]$.

	LO	NLO	N ² LO	N ³ LO	N ⁴ LO	Total
$\Gamma^{O(\alpha_s^6)} _{\text{Conv.}}$	218.66 ^{-40.95} _{+57.39}	116.62 ^{+2.08} _{-15.59}	14.44 ^{+25.23} _{-41.60}	-8.70 ^{+13.84} _{-8.02}	-3.57 ^{+1.74} _{+6.64}	337.45 ^{+1.94} _{-1.18}
$\Gamma^{O(\alpha_s^6)} _{\text{PMC}_\infty}$	299.57	95.22	-41.82	-19.59	3.04	336.42
$\Gamma^{O(\alpha_s^6)} _{\text{PMCm}}$	289.83 ^{+0.96} _{-0.24}	91.19 ^{-0.51} _{+0.29}	-31.78 ^{+0.30} _{-0.11}	-13.36	1.92	337.80 ^{+0.75} _{-0.06}

$$\Gamma(H \rightarrow gg)|_{\text{Conv.}} = 337.45^{+6.27+1.21+0.02}_{-6.20-1.21-0.02} \text{ keV}, \quad (12)$$

$$\Gamma(H \rightarrow gg)|_{\text{Conv.}} = 337.45^{+6.67}_{-6.43} \text{ keV}, \quad (14)$$

$$\Gamma(H \rightarrow gg)|_{\text{PMC}_\infty} = 336.42^{+6.21+1.22+0.03}_{-6.14-1.20-0.01} \text{ keV}, \quad (13)$$

$$\Gamma(H \rightarrow gg)|_{\text{PMC}_\infty} = 336.42^{+6.33}_{-6.26} \text{ keV}, \quad (15)$$

where the errors are $\Delta\alpha_s(M_Z) = \pm 0.0009$ (which leads to $\Lambda_{\text{mMOM}}^{n_f=5} = 362.0^{+36.6}_{-18.0}$ MeV), $\Delta M_H = \pm 0.17$ GeV, and $\Delta M_t = \pm 0.7$ GeV, respectively. Here, the conventional predictions are achieved by fixing $\mu_r = M_H$.

Using the PMC_∞ approach, the PMC scales at each order are no longer evaluated as a perturbative series, thus avoiding the *first type of residual scale dependence*. As mentioned above, all the PMC_∞ scales are around $M_H \exp(-5/6) \sim 54$ GeV, so the *second type of residual scale dependence* is small owing to the convergent behavior at higher orders. As a further step of making a conservative estimation on the contributions from the uncalculated $n+1$ th-order terms, we set its PMC_∞ scale μ_n to be within the region of the latest determined PMC_∞ scale μ_{n-1} , e.g., $\mu_n \in [\mu_{n-1}/2, 2\mu_{n-1}]$, and set $\Delta\Gamma^{O(\alpha_s^{i+1})}|_{\text{PMC}_\infty} = \pm \frac{M_i G_F}{36\sqrt{2}\pi} |C_{i,\text{Conf}} a_s^{i+2}(\mu_{i+1})|_{\text{MAX}}$ with $i = I, II, III, IV, V$, respectively. Numerically, we obtained $\Delta\Gamma^{O(\alpha_s^2)}|_{\text{PMC}_\infty} = \pm 2.99$ keV, $\Delta\Gamma^{O(\alpha_s^3)}|_{\text{PMC}_\infty} = \pm 1.80$ keV, $\Delta\Gamma^{O(\alpha_s^4)}|_{\text{PMC}_\infty} = \pm 1.25$ keV, $\Delta\Gamma^{O(\alpha_s^5)}|_{\text{PMC}_\infty} = \pm 0.47$ keV, and $\Delta\Gamma^{O(\alpha_s^6)}|_{\text{PMC}_\infty} = \pm 0.09$ keV. Note that the estimated errors may underestimate the contributions listed in Table 1, which need to be multiplied by {57, 27, 6, 5}, respectively. Thus, the numerical results for $\Delta\Gamma^{O(\alpha_s^i)}|_{\text{PMC}_\infty}$ need to be multiplied by 5 as analogy. Similarly, one can obtain the corresponding values for the conventional scale-setting approach, which are ± 11.98 keV, ± 4.60 keV, ± 1.40 keV, ± 0.73 keV, and ± 0.14 keV, respectively. To match the center values shown in Table 1, these values need to be multiplied by {10, 3, 6, 5}. Moreover, the numerical results for $\Delta\Gamma^{O(\alpha_s^6)}|_{\text{Conv.}}$ need to be multiplied by 5 as analogy. Those values are slightly larger than those of PMC_∞ owing to a larger perturbative coefficient C_k than the conform coefficient at each order, even if it is compensated by a smaller α_s value.

As a summary, we have presented a detailed analysis of the Higgs-boson decay $H \rightarrow gg$ up to α_s^6 -order, and we obtain

where the errors are squared averages of those from $\Delta\alpha_s(M_Z)$, ΔM_H , ΔM_t , and the uncertainty of the renormalization scale within the region of $[M_H/2, 2M_H]$. The errors are dominated by $\Delta\alpha_s(M_Z)$, followed by the choice of renormalization scale and accuracy of Higgs mass. If the value of $\alpha_s(M_Z)$ can be measured accurately to avoid the error from $\Delta\alpha_s(M_Z)$, we will obtain

$$\Gamma(H \rightarrow gg)|_{\text{Conv.}} = 337.45^{+2.29}_{-1.69} \text{ keV}, \quad (16)$$

$$\Gamma(H \rightarrow gg)|_{\text{PMC}_\infty} = 336.42^{+1.22}_{-1.20} \text{ keV}. \quad (17)$$

The Higgs-boson decay $H \rightarrow gg$ provides another successful example for the application of the PMC_∞ scale-setting method to high-energy processes. Up to N⁴LO QCD corrections, the pQCD predictions under PMC_∞ and conventional scale-setting approaches are consistent with each other. However, the conventional renormalization scale uncertainties are still sizable, i.e., approximately 1% when varying the renormalization scale μ_r within the range of $[\frac{1}{2}M_H, 2M_H]$. By applying PMC_∞, the α_s values at lower orders are definitely fixed by the requirement of intrinsic conformality, the conventional renormalization scale ambiguity is eliminated, and the residual scale dependence from the original PMC multi-scale-setting approach is also highly suppressed. Thus, a more precise test of the SM can be achieved.

APPENDIX: CONFORMAL COEFFICIENTS AND PMC_∞ SCALES UP TO α_s^6 -ORDER LEVEL

Applying the PMC_∞ scale-setting approach together with the general "degeneracy" pattern of the QCD theory [36], the perturbative series of the decay width $\Gamma(H \rightarrow gg)$ under the mMOM-scheme is

$$\begin{aligned}
 \Gamma(H \rightarrow gg) = & \frac{M_H^3 G_F}{36 \sqrt{2} \pi} \left[A_{\text{Conf}} a_s^2(\mu_r) + (B_{\text{Conf}} + 2A_{\text{Conf}} B_{\beta_0} \beta_0) a_s^3(\mu_r) + \left(C_{\text{Conf}} + 3B_{\text{Conf}} C_{\beta_0} \beta_0 + 3A_{\text{Conf}} B_{\beta_0}^2 \beta_0^2 \right. \right. \\
 & + 2A_{\text{Conf}} B_{\beta_0} \beta_1 \left. \right) a_s^4(\mu_r) + \left(D_{\text{Conf}} + 7A_{\text{Conf}} B_{\beta_0}^2 \beta_1 \beta_0 + 4C_{\text{Conf}} D_{\beta_0} \beta_0 + 6B_{\text{Conf}} C_{\beta_0}^2 \beta_0^2 + 4A_{\text{Conf}} B_{\beta_0}^3 \beta_0^3 \right. \\
 & + 2A_{\text{Conf}} B_{\beta_0} \beta_2 + 3B_{\text{Conf}} C_{\beta_0} \beta_1 \left. \right) a_s^5(\mu_r) + \left(E_{\text{Conf}} + 8A_{\text{Conf}} B_{\beta_0}^2 \beta_2 \beta_0 + \frac{27}{2} B_{\text{Conf}} C_{\beta_0}^2 \beta_1 \beta_0 + 5D_{\text{Conf}} E_{\beta_0} \beta_0 \right. \\
 & + \frac{47}{3} A_{\text{Conf}} B_{\beta_0}^3 \beta_1 \beta_0^2 + 10C_{\text{Conf}} D_{\beta_0}^2 \beta_0^2 + 10B_{\text{Conf}} C_{\beta_0}^3 \beta_0^3 + 5A_{\text{Conf}} B_{\beta_0}^4 \beta_0^4 + 2A_{\text{Conf}} B_{\beta_0} \beta_3 + 3B_{\text{Conf}} C_{\beta_0} \beta_2 \\
 & \left. \left. + 4A_{\text{Conf}} B_{\beta_0}^2 \beta_1^2 + 4C_{\text{Conf}} D_{\beta_0} \beta_1 \right) a_s^6(\mu_r) \right] + O[a_s^7(\mu_r)]. \tag{A1}
 \end{aligned}$$

To compare Eq. (1) with Eq. (18), one can systematically determine the conformal coefficients and PMC_∞ scales for $\Gamma(H \rightarrow gg)$ up to the α_s^6 -order level as follows:

$$A_{\text{Conf}} = C_0, \tag{A2}$$

$$B_{\text{Conf}} = C_1 \left(n_f = \frac{33}{2} \right), \tag{A3}$$

$$C_{\text{Conf}} = C_2 \left(n_f = \frac{33}{2} \right) - 2A_{\text{Conf}} B_{\beta_0} \bar{\beta}_1, \tag{A4}$$

$$D_{\text{Conf}} = C_3 \left(n_f = \frac{33}{2} \right) - 2A_{\text{Conf}} B_{\beta_0} \bar{\beta}_2 - 3B_{\text{Conf}} C_{\beta_0} \bar{\beta}_1, \tag{A5}$$

$$\begin{aligned}
 E_{\text{Conf}} = & C_4 \left(n_f = \frac{33}{2} \right) - 2A_{\text{Conf}} B_{\beta_0} \bar{\beta}_3 - 3B_{\text{Conf}} C_{\beta_0} \bar{\beta}_2 \\
 & - 4A_{\text{Conf}} B_{\beta_0}^2 \bar{\beta}_1^2 - 4C_{\text{Conf}} D_{\beta_0} \bar{\beta}_1
 \end{aligned} \tag{A6}$$

and

$$\ln \frac{\mu_r^2}{\mu_1^2} = \frac{C_1 - B_{\text{Conf}}}{2A_{\text{Conf}} \beta_0}, \tag{A7}$$

$$\ln \frac{\mu_r^2}{\mu_{\text{II}}^2} = \frac{C_2 - C_{\text{Conf}} - 3A_{\text{Conf}} B_{\beta_0}^2 \beta_0^2 - 2A_{\text{Conf}} B_{\beta_0} \beta_1}{3B_{\text{Conf}} \beta_0}, \tag{A8}$$

$$\ln \frac{\mu_r^2}{\mu_{\text{III}}^2} = \frac{C_3 - D_{\text{Conf}} - 7A_{\text{Conf}} B_{\beta_0}^2 \beta_1 \beta_0 - 6B_{\text{Conf}} C_{\beta_0}^2 \beta_0^2 - 4A_{\text{Conf}} B_{\beta_0}^3 \beta_0^3 - 2A_{\text{Conf}} B_{\beta_0} \beta_2 - 3B_{\text{Conf}} C_{\beta_0} \beta_1}{4C_{\text{Conf}} \beta_0}, \tag{A9}$$

$$\begin{aligned}
 \ln \frac{\mu_r^2}{\mu_{\text{IV}}^2} = & (C_4 - E_{\text{Conf}} - 8A_{\text{Conf}} B_{\beta_0}^2 \beta_2 \beta_0 - \frac{27}{2} B_{\text{Conf}} C_{\beta_0}^2 \beta_1 \beta_0 - \frac{47}{3} A_{\text{Conf}} B_{\beta_0}^3 \beta_1 \beta_0^2 - 10C_{\text{Conf}} D_{\beta_0}^2 \beta_0^2 - 10B_{\text{Conf}} C_{\beta_0}^3 \beta_0^3 \\
 & - 5A_{\text{Conf}} B_{\beta_0}^4 \beta_0^4 - 2A_{\text{Conf}} B_{\beta_0} \beta_3 - 3B_{\text{Conf}} C_{\beta_0} \beta_2 - 4A_{\text{Conf}} B_{\beta_0}^2 \beta_1^2 - 4C_{\text{Conf}} D_{\beta_0} \beta_1) / 5D_{\text{Conf}} \beta_0,
 \end{aligned} \tag{A10}$$

where $\bar{\beta}_1 = \beta_1(n_f = \frac{33}{2}) = -107$, $\bar{\beta}_2 = \beta_2(n_f = \frac{33}{2}) = -2001.29$, and C_k are the perturbative coefficients.

References

- [1] A. Djouadi, *Phys. Rept.* **457**, 1 (2008)
- [2] T. Inami, T. Kubota, and Y. Okada, *Z. Phys. C* **18**, 69 (1983)
- [3] A. Djouadi, M. Spira, and P. M. Zerwas, *Phys. Lett. B* **264**, 440 (1991)
- [4] D. Graudenz, M. Spira, and P. M. Zerwas, *Phys. Rev. Lett.* **70**, 1372 (1993)
- [5] S. Dawson and R. Kauffman, *Phys. Rev. D* **49**, 2298 (1994)
- [6] M. Spira, A. Djouadi, D. Graudenz *et al.*, *Nucl. Phys. B* **453**, 17 (1995)
- [7] S. Dawson and R. P. Kauffman, *Phys. Rev. Lett.* **68**, 2273 (1992)
- [8] K. G. Chetyrkin, B. A. Kniehl, and M. Steinhauser, *Phys. Rev. Lett.* **79**, 353 (1997)
- [9] K. G. Chetyrkin, B. A. Kniehl, and M. Steinhauser, *Nucl. Phys. B* **510**, 61 (1998)
- [10] P. A. Baikov and K. G. Chetyrkin, *Phys. Rev. Lett.* **97**, 061803 (2006)
- [11] F. Herzog, B. Ruijl, T. Ueda *et al.*, *JHEP* **1708**, 113 (2017)
- [12] S. J. Brodsky and X. G. Wu, *Phys. Rev. D* **85**, 034038 (2012)

- [13] S. J. Brodsky and X. G. Wu, *Phys. Rev. Lett.* **109**, 042002 (2012)
- [14] M. Mojaza, S. J. Brodsky, and X. G. Wu, *Phys. Rev. Lett.* **110**, 192001 (2013)
- [15] S. J. Brodsky, M. Mojaza, and X. G. Wu, *Phys. Rev. D* **89**, 014027 (2014)
- [16] X. G. Wu, S. J. Brodsky and M. Mojaza, *Prog. Part. Nucl. Phys.* **72**, 44 (2013)
- [17] X. G. Wu, Y. Ma, S. Q. Wang *et al.*, *Rep. Prog. Phys.* **78**, 126201 (2015)
- [18] M. Gell-Mann and F. E. Low, *Phys. Rev.* **95**, 1300 (1954)
- [19] S. J. Brodsky and P. Huet, *Phys. Lett. B* **417**, 145 (1998)
- [20] S. Q. Wang, X. G. Wu, X. C. Zheng *et al.*, *Eur. Phys. J. C* **74**, 2825 (2014)
- [21] D. M. Zeng, S. Q. Wang, X. G. Wu *et al.*, *J. Phys. G* **43**, 075001 (2016)
- [22] J. Zeng, X. G. Wu, S. Bu *et al.*, *J. Phys. G* **45**, 085004 (2018)
- [23] X. C. Zheng, X. G. Wu, S. Q. Wang *et al.*, *JHEP* **10**, 117 (2013)
- [24] X. G. Wu, J. M. Shen, B. L. Du *et al.*, *Prog. Part. Nucl. Phys.* **108**, 103706 (2019)
- [25] L. Di Giustino, S. J. Brodsky, S. Q. Wang *et al.*, *Phys. Rev. D* **102**, 014015 (2020)
- [26] W. Celmaster and R. J. Gonsalves, *Phys. Rev. D* **20**, 1420 (1979)
- [27] W. Celmaster and R. J. Gonsalves, *Phys. Rev. Lett.* **42**, 1435 (1979)
- [28] W. Celmaster and R. J. Gonsalves, *Phys. Rev. Lett.* **44**, 560 (1980)
- [29] W. Celmaster and R. J. Gonsalves, *Phys. Rev. D* **21**, 3112 (1980)
- [30] J. A. Gracey, *J. Phys. A* **46**, 225403 (2013)
- [31] L. von Smekal, K. Maltman, and A. Sternbeck, *Phys. Lett. B* **336**, (2009)
- [32] J. Zeng, X. G. Wu, X. C. Zheng *et al.*, *Chin. Phys. C* **44**, 113102 (2020)
- [33] S. J. Brodsky and H. J. Lu, *Phys. Rev. D* **51**, 3652 (1995)
- [34] B. Ruijl, T. Ueda, J. A. M. Vermaseren *et al.*, *JHEP* **1706**, 040 (2017)
- [35] R.L. Workman *et al.* (Particle Data Group), *PTEP* **2022**, 083C01 (2022)
- [36] H. Y. Bi, X. G. Wu, Y. Ma, *et al.*, *Phys. Lett. B* **748**, 13 (2015)